

## AN ANALYSIS OF HYDATID CYST SURGERIES IN TEHRAN MILAD HOSPITAL, IRAN, DURING 2001- 2004

A. Pezeshki<sup>1</sup>, E.B. Kia<sup>2</sup>, A. Gholizadeh<sup>3</sup>, A.Koohzare<sup>4</sup>

### ABSTRACT

Hydatidosis is a zoonotic disease which is due to infectivity with larval stage of dog tapeworm, "*Echinococcus granulosus*". The disease is chronic and cysts can be lodged in different organs of the intermediate hosts. It has cosmopolitan distribution and impact health and economical challenges for the many countries throughout the world. In Iran, human cases are constantly reported from different medical centers, Therefore, accurate information on the distribution of the disease is first step for the control and prevention. In this descriptive study, demographic information ( sex, age, occupation ) about patients who underwent hydatid cyst surgeries during 2001-2004 in Tehran Milad Hospital were collected and analyzed.

Among 78 patients who had hydatid cyst operations in Milad hospital, 56.5% were female and 43.5% male. Liver was the most commonly involved organ. According to the result of this study, females were found more infected with hydatid cysts than males.

**KEY WORDS:** Hydatid cyst, surgery, Iran.

Pak J Med Sci January - March 2007 Vol. 23 No. 1 138-140

### INTRODUCTION

Hydatid disease is a zoonotic infection caused by *Echinococcus* species and is divided into two histopathologic forms: cystic and alveolar form. The cystic form of the disease is more common and usually characterized with large, fluid – filled, unilocular cysts in which *Echinococcus granulosus* is the causative agent.<sup>1</sup> Hydatid cyst is found in human and some other mammalian tissues, especially liver and lungs and

causes echinococcosis.<sup>2</sup> The disease poses an important public health problem in many areas of the world, particularly among populations that practice sheep husbandry.<sup>3</sup> Cystic echinococcus is endemic in Iran, and is maintained in three distinct cycles, a livestock / dog domestic cycle, a desert cycle between dogs and camels, and a sylvatic cycle between wild carnivores and wild ruminants.<sup>4</sup>

The *E. granulosus* cycle is typically a dog-sheep cycle and implies contamination of a sheep through the feces of an infected dog. Humans accidentally take the place of the sheep in the parasite cycle through close contact with an infected dog.<sup>5</sup> Humans are usually a "dead- end" for the parasite.<sup>6</sup> Serological methods, used for the diagnosis of hydatid disease are direct hemagglutination, latex agglutination, immunoelectrophoresis, skin tests and ELISA.<sup>7</sup> Radiological imaging methods such as computed tomography (CT) and MR imaging play an important role in the diagnosis.<sup>8,9</sup>

Therapy for echinococcosis is based on the size, location, and symptomatic complications

1. Dr. A. Pezeshki
  2. Dr. E.B. Kia
  3. Dr. A. Gholizadeh
  4. Dr. A.Koohzare,  
Tehran Milad Hospital,  
Tehran, Iran.
- 1-3: Department of Medical Parasitology & Mycology,  
School of Public Health &  
Institute of Health Research,  
Tehran University of Medical Sciences,  
P.O.Box : 14155 - 6446,  
Tehran, Iran.

Correspondences:

Dr. A. Pezeshki  
E-Mail: pezeshkii@yahoo.com

\* Received for Publication: March 2, 2006

\* Accepted: September 30, 2006

of the cysts. Surgery is the treatment of choice.<sup>10-12</sup> Preventive strategies would, therefore, involve health education, reduced contacts with dogs and sheep and with effective disposal of their wastes.<sup>13</sup>

In Iran, human cases are constantly reported from different medical centers. Milad hospital is located in Tehran, capital city, and people from different parts of the country are referred here for hydatid surgery. In this retrospective study the hydatid surgeries in this hospital during five years were analyzed.

### PATIENTS AND METHODS

Milad hospital is a private sector healthcare facility in Tehran which attract patients from all parts of the country who are referred here for hydatid surgery. In this descriptive study, demographic information (sex, age, occupation) about patients who underwent hydatid cyst surgeries during 2001-2004 in Tehran Milad Hospital were collected from hospital files and analyzed.

### RESULTS

A total of seventy eight patients 34 (43.5%) male and 44 (56.5%) female were enrolled in this study. Majority of the patients 31.5% were in the group of 32-40 years.(Table-I) Liver was the most common organ involved i.e (69.23%). The involvement of other organs was lung (11.53%), spleen (7.69%), abdomen (3.84%), spinal cord (2.56%), under diaphragm (1.28%), peritoneum (1.28%), kidney (1.28%) and pancreas (1.28%) (Table-II).

Table-I: Age of patients infected with hydatid cyst

Age	Number of patients
14-22	11
23-31	8
32-40	17
41-49	11
50-58	14
59-67	4
68-76	13

Table-II: Involvement of different organ (n=78)

Affected organ	Number of patients	(%)	Males	Females
Liver	54	(69.23)	20	34
Lung	9	(11.53)	5	4
Spleen	6	(7.69)	4	2
Abdomen	3	(3.84)	2	1
Spinal cord	2	(2.56)	1	1
Under diaphragm	1	(1.28)	1	-
Peritoneum	1	(1.28)	1	-
Kidney	1	(1.28)	-	1
Pancreas	1	(1.28)	-	1

### DISCUSSION

Hydatid disease is a medical and public health problem, especially in endemic parts of the world. The transmission of the disease mostly depends on the close relationship between human beings and canine carnivores. Humans are intermediate hosts and are being exposed to the parasite by fecal-oral and hand-to-mouth spread ways.<sup>1</sup> Iran is an important endemic focus of cystic hydatid disease.<sup>2</sup> Investigations on echinococcosis, which have been performed in this country showed the presence of infection among human.<sup>14-19</sup> According to the result of this study, females were found more infected with hydatid cysts than males. Different organs are involved with cyst, however, liver is the most affected organ. During a study carried out by Fattahi et al. between 1991-1997.<sup>20</sup> In Markazi province women had more hydatid surgeries than men and the most affected organ was liver. Nilforoushan et al<sup>21</sup> In Isfahan and Moradi & Majidpoor<sup>22</sup> in Kordestan reported similar results. However, Adibpoor et al.<sup>23</sup> in Tabriz found more hydatid surgeries in men than women and lung was the most affected organ. In conclusion, echinococcosis is still an important health problem in Iran that needs further studies. Therefore, accurate information on the distribution of the disease is first step for the control and prevention. Moreover, it is necessary that in each province the role of different intermediate hosts and the strains of *E. granulosus* in human and animals be investigated.

## REFERENCES

- Smith DM, Rickman LS. Echinococcosis: A drop of water – review on human disease, diagnosis, and management. *Infect Disease Clin Practice* 2001;10:355-9.
- Harandi MF, Hobbs RP, Adams PJ, Mobedi I, Morgan – Ryan UM, Thompson RC. Molecular and morphological characterization of *Echinococcus granulosus* of human and animal origin in Iran. *Parasitol* 2002;125(4):367-73.
- Anonymous. Guidelines for surveillance, prevention and control of echinococcosis / hydatidosis. WHO/FAO/UNEP/8128. 1981.
- Dalimi AGH, Motamedi M, Hosseini B, Mohammadian H, Malaki Z, Ghamari, et al. Echinococcosis / hydatidosis in western Iran. *Vet. Parasitol* 2002;105:161-71.
- Horchani AY, Nouria, Kabier I. Hydatid cyst of the kidney: A report of 147 controlled cases. *Eur Urol* 2000;33:461-7.
- Macpherson CNL. An active intermediate host role for man in the life cycle of *Echinococcus granulosus* in Turkana, Kenya. *Am J Trop Med Parasitol* 1983;31:397-404.
- Romos G, Orduna A, Garcia-yuste M. Hydatid cyst of the lung: Diagnosis and treatment. *World J Surg* 2001;25:46-57.
- Tuzun M, Hekimoglu B. Various locations of cystic and alveolar hydatid disease. CT appearances. *J Comput Assist Tomogr* 2001; 25: 81-7.
- Akhan O, Ozmen MN, Dincer A, Sayek I, Gocmen A. Liver hydatid disease: long-term results of percutaneous treatment. *Radiology* 1996;198:259-64.
- Kammerer WS, Schantz PM. Echinococcal disease. *Infect Dis Clin North Am* 1993;7:605-18.
- Ammann RW, Eckert J, Cestods. *Echinococcus* Gastro. *Clin North Am* 1996;25:655-89.
- Garcia LS, Bruckner DA. Tissue cestodes: larval form. In: *Diagnostic medical Parasitology*. 4ed. Washington, DC. ASM press 2001;387-403.
- Palmer SR, Biffin AH, Crig PS, Walters TM. Control of hydatid disease in Wales. *BMJ* 1993;12:674-5.
- Bastani B, Dehdashti F. Hepatic hydatid disease in Iran with review of the literature. *Mt Sinai J Med* 1995;62-9.
- Kelsey DS, Shaffner LS. Pulmonary echinococcosis: the geographic history scores again. *NC Med J* 1984; 45 (4):218-19.
- Panahi F. Costal echinococcosis. Report of a case and review of the literature. *Sem. Hop* 1978;8-15(43-44):1389-92.
- Lomheh F. Immunofluorescence in the serodiagnosis of human hydatidosis. *Acta Med Iran* 1997;20(1-2):27-35.
- Sabbaghian HN, Hoghochi, Ghaderian E. A survey on the prevalence of echinococcosis in shahrkord Iran. *Bull Soc Pathol Exot Filials*, 1975;68(6):547.
- Nasseh GA, Khadivi B. Epidemiological and clinical aspects of echinococcosis in east Iran. *J Trop Med Hyg.* 1975;78(6):120-22.
- Fattahi F, Davami M, Mosayyebi M, Study on hydatid cysts surgeries in Markazi province, Iran. *Proceedings of 3rd National Congress of Medical Parasitology, IRAN*, 27 Feb.-1 Mar. 2001; SARI, IRAN.
- Nilforoushan MR, Davoodi VS, Kia I EB. Mobedi, Study on Hydatid cysts surgeries in Isfahan Educational centers during 1996-2001. 13th Iranian congress on Infectious Disease and Tropical / Medicine 11-15 Dec, 2004.
- Moradi G, Majidpoor MS. Epidemiological study on Hydatid cysts in Kordestan province, Iran. during 2000-2004. 13th Iranian congress on Infectious Disease and Tropical / Medicine 11-15 Dec. 2004.
- Adibpoor MR, Jamali, Sarkarty F. Survey on hydatid cysts surgeries in Tabriz (Iran) Hospitals during 1991-1997. *Proceedings of 3rd National Congress of Medical Parasitology, IRAN*, 27 Feb. 1Mar. 2001; SARI, IRAN.